

References

1. Moscicki EK. Identification of suicide risk factors using epidemiologic studies. *Psychiatr Clin North Am* 1997;20:499-517.
2. Moscicki EK. Epidemiology of suicidal behavior. *Suicide Life Threat Behav* 1995;25:22-35.
3. Barraclough BM, Pallis DJ. Depression followed by suicide: a comparison of depressed suicides with living depressives. *Psychol Med* 1975;5:55-61.
4. Henriksson MM, Aro HM, Marttunen MJ, Heikkinen ME, Isometsä ET, Kuopasalmi KI, and others. Mental disorders and comorbidity in suicide. *Am J Psychiatry* 1993;150:935-40.
5. Isometsä ET, Henriksson MM, Aro HM, Heikkinen ME, Kuopasalmi KI, Lönnqvist JK. Suicide in major depression. *Am J Psychiatry* 1994;151:530-6.
6. Modestin J, Kopp W. Study on suicide in depressed patients. *J Affect Disord* 1988;15:157-62.
7. Keller MB, Hanks DL. Anxiety symptom relief in depression treatment outcomes. *J Clin Psychiatry* 1995;56(Suppl 6):22-9.
8. Lecrubier Y. The impact of comorbidity on the treatment of panic disorder. *J Clin Psychiatry* 1998;59(Suppl 8):11-4.
9. Sherbourne CD, Wells KB. Course of depression in patients with comorbid anxiety disorders. *J Affect Disord* 1997;43:245-50.
10. Fawcett J, Scheftner WA, Fogg L, Clark DC, Young MA, Hedeker D, and others. Time-related predictors of suicide in major affective disorder. *Am J Psychiatry* 1990;147:1189-94.
11. Johnson J, Weissman MM, Klerman GL. Panic disorder, comorbidity, and suicide attempts. *Arch Gen Psychiatry* 1990;47:805-8.
12. Vollrath M, Angst J. Outcome of panic and depression in a seven-year follow-up: results of the Zurich study. *Acta Psychiatr Scand* 1989;80:591-6.
13. Joffe RT, Bagby RM, Levitt A. Anxious and nonanxious depression. *Am J Psychiatry* 1993;150:1257-8.
14. Canetto SS, Sakinofsky I. The gender paradox in suicide. *Suicide Life Threat Behav* 1998;28:1-23.
15. Rich CL, Ricketts JE, Fowler RC, Young D. Some differences between men and women who commit suicide. *Am J Psychiatry* 1988;145:718-22.
16. Spitzer RL, Williams JBW, Gibbon M, First M. Structured Clinical Interview for DSM-IV. Washington (DC): American Psychiatric Press; 1995.

Clinical Implications

- Further evidence is provided regarding possible risk factors for suicidality in patients with major depression.
- No interaction between anxiety and sex on suicidality was found.
- The high frequency of suicidality in patients with major depression is reconfirmed.

Limitations

- This study used a retrospective design.
- Subjects were recruited from a specialized depression clinic.
- Only cross-sectional data were used to determine suicidality.

17. Endicott J, Spitzer RL. A diagnostic interview: the schedule for affective disorders and schizophrenia. *Arch Gen Psychiatry* 1978;35:837-44.
18. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56-62.
19. Beck AT, Ward CH, Mendelson M, and others. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561-71.
20. Ohring R, Apter A, Ratzoni G, Weizman R, Tyano S, Plutchik R. State and trait anxiety in adolescent suicide attempters. *J Am Acad Child Adolesc Psychiatry* 1996;35:154-7.
21. Bell CJ, Nutt DJ. Serotonin and panic. *Br J Psychiatry* 1998;172:465-71.
22. Greist JH. Anxiety disorders: the role of serotonin. *J Clin Psychiatry* 1996;57(Suppl 6):3-4.
23. Nordstrom P, Asberg M. Suicide risk and serotonin. *Int Clin Psychopharmacol* 1992;6(Suppl 6):12-21.
24. Brown GR, Anderson B. Psychiatric morbidity in adult inpatients with childhood histories of sexual and physical abuse. *Am J Psychiatry* 1991;148:55-61.
25. Bryer JB, Nelson BA, Miller JB, Krol PA. Childhood sexual and physical abuse as factors in adult psychiatric illness. *Am J Psychiatry* 1987;144:1426-30.