



Tips for Survivors:

COPING WITH GRIEF AFTER A DISASTER OR TRAUMATIC EVENT

Grief is the normal response of sorrow, heartache, and confusion that comes from losing someone or something important to you. Grief can also be a common human response after a disaster or other traumatic event.

This tip sheet contains information about grief, the grieving process, and what happens when the process is interrupted and complicated or traumatic grief occurs. It also offers tips and resources for coping with both types of grief.

Grief is a part of life. It is a strong, sometimes overwhelming reaction to death, divorce, job loss, a move, or loss of health due to illness. It can also occur after disasters or other traumatic events.



What Is Grief?

Grief is a part of life. It is a strong, sometimes overwhelming reaction to death, divorce, job loss, a move, or loss of health due to illness. It can also occur after disasters or other traumatic events. If you are experiencing grief, you may feel empty and numb, or unable to feel joy or sadness. You may also feel angry. You may experience physical reactions, including the following:

- Trembling or shakiness
- Muscle weakness
- Nausea and trouble eating
- Trouble sleeping or difficulty breathing
- Dry mouth

While grieving, you may have nightmares, withdraw socially, and not want to participate in your usual activities.

Eventually, you may also notice positive changes in yourself from your experience of loss, such as the following:

- Becoming more understanding and tolerant
- Having increased appreciation for relationships and loved ones
- Feeling grateful for the presence of those in your community who are loving and caring
- Experiencing enhanced spiritual connection
- Becoming more socially active
- Getting involved in advocacy to help people who have experienced similar losses

How Long Does Grief Usually Last?

Grief is different for every person and every loss. The manner of the death or loss and your personal circumstances can affect how long you experience grief.

If you are experiencing grief now, it may be difficult to imagine an end to the pain you are feeling. Even though your life may not be the same as it was before, the grief will eventually ease, and the grieving process will allow you to accept the loss and move forward with your life.




Coping With Grief

You can take steps to help yourself cope with grief. Talking to others who understand and respect how you feel—family members, faith leaders, people you trust—is a helpful way to cope with your grief. Recognize that although you might still have these feelings over a long period, they will likely be less intense over time. Try to take care of yourself physically by exercising, eating healthy meals, and getting enough sleep. Allow yourself to feel joy and sadness, and to cry when you need to.

Going through the grieving process and healing from loss does not mean forgetting about loved ones who have died. You may still feel deeply connected to the person you have lost, but you will also be able to imagine a life without him or her. You will start to be reengaged in daily life and reconnected to others. When moving out of grief, you may start to experience the following:

- Feeling the pain of your loss without resisting or avoiding it
- Adjusting to a new reality in which your loved one is no longer present
- Forming new relationships



After a disaster or other traumatic event, many factors can disrupt the normal process of grieving, raising the risk for complicated or traumatic grief.


If you are not experiencing these changes and adjustments after many months, and you feel “stuck” in your deep and overwhelming feelings of loss, then you may be experiencing what is known as complicated or traumatic grief.

What Is Complicated or Traumatic Grief?

After a disaster or other traumatic event, many factors can disrupt the normal process of grieving, raising the risk for complicated or traumatic grief. Whether a disaster is natural or human-caused, grief at the loss of loved ones may be compounded by sorrow and anger at the loss of home and possessions, as well as fear or anxiety about the loss of a sense of safety and security in the community. In some cases, meeting immediate physical needs after a disaster may take priority over grieving the loss of life, which can delay, prolong, or complicate the normal grieving process.

Complicated or traumatic grief is grief that does not end and does not help you make progress toward getting back to your usual activities and routine. For most people, intense feelings of


grief will lessen gradually over time, beginning to ease within 6 months of the loss. But if you’re experiencing complicated or traumatic grief, you may not feel any reduction of grief over many months or even years. Your feelings of sadness, anger, and loneliness may even become more intense over time.



These are some of the symptoms of complicated or traumatic grief:

- Feeling deeply angry about the death or loss
- Being unable to think about anything but your loved one
- Not wanting any reminders of your loved one at all
- Having nightmares or intrusive thoughts
- Feeling deep loneliness and longing for the person you lost
- Feeling distrustful of others
- Feeling unable to maintain regular activities or fulfill responsibilities
- Feeling bitterness about life and envying others not affected by grief
- Being unable to enjoy life or remember happy times with your loved one

If you are experiencing any of these symptoms, you may need to seek help in order to move forward and begin the healing process. You can find help by using the resources listed on this page, talking to your doctor, or seeing another health care professional.



Helpful Resources

SAMHSA Disaster Technical Assistance Center

Toll-free: 1-800-308-3515

Website: <https://www.samhsa.gov/dtac>

SAMHSA Behavioral Health Disaster Response Mobile App

Website: <https://store.samhsa.gov/product/PEP13-DKAPP-1>

Administration for Children and Families

Website: <https://www.acf.hhs.gov>

Mental Health America*

Website: <http://www.mentalhealthamerica.net>

National Alliance on Mental Illness (NAMI)*

Toll-free helpline: 1-800-950-NAMI (1-800-950-6264)

Website: <https://www.nami.org>

Treatment Locators

Behavioral Health Treatment Facility Locator

Toll-free: 1-800-662-HELP (1-800-662-4357) (24/7 English and español)

TDD: 1-800-487-4889

Website: <https://www.findtreatment.samhsa.gov>

MentalHealth.gov

Website: <https://www.mentalhealth.gov>

MentalHealth.gov provides U.S. government information and resources on mental health.

Helplines

SAMHSA Disaster Distress Helpline

Website: <https://www.samhsa.gov/find-help/disaster-distress-helpline>

Call 1-800-985-5990 or text "TalkWithUs" to 66746 to get help and support 24/7.

National Suicide Prevention Lifeline

Toll-free: 1-800-273-TALK (1-800-273-8255)

TTY: 1-800-799-4TTY (1-800-799-4889)

Website: <https://www.samhsa.gov>

This resource can be found by accessing the National Suicide Prevention Lifeline box on the SAMHSA website.

*Note: The views, opinions, and content expressed in this publication do not necessarily reflect the views, opinions, or policies of the Center for Mental Health Services (CMHS), the Substance Abuse and Mental Health Services Administration (SAMHSA), or the U.S. Department of Health and Human Services (HHS).



Complicated grief therapy as a new treatment approach

Julie Loebach Wetherell, PhD



Complicated grief therapy (CGT) is a relatively new psychotherapy model designed to address symptoms of complicated grief. Drawn from attachment theory and with roots in both interpersonal therapy (IPT) and cognitive-behavioral therapy, CGT includes techniques similar to prolonged exposure (repeatedly telling the story of the death and in vivo exposure activities). The treatment also involves focusing on personal goals and relationships. CGT has been demonstrated to be effective in a trial in which participants with complicated grief were randomly assigned to CGT or IPT; individuals receiving CGT responded more quickly and were more likely to respond overall (51% vs 28%). This article briefly summarizes the conceptual underpinnings of CGT, discusses the empirical evidence for its efficacy, describes its techniques, and presents a case example of a client treated in a 16-session manualized CGT protocol. The article concludes with a description of future research directions for CGT.

© 2012, LLS SAS

Dialogues Clin Neurosci. 2012;14:159-166.

Keywords: *bereavement; traumatic grief; treatment; psychotherapy; cognitive behavior therapy*

Author affiliations: Psychology Service, VA San Diego Healthcare System and Department of Psychiatry, University of California, San Diego, California, USA

Introduction

Facilitating recovery from loss has been a staple of psychotherapy since long before the entity known variously as “complicated grief,” “traumatic grief,” “complicated bereavement,” “prolonged grief disorder,” or “pathologic grief” was identified as a form of suffering distinct from normal bereavement or depression. Clinicians have described numerous forms of treatment for bereavement-related distress, relying on different conceptualizations of the problem and different therapeutic techniques,^{1,2} including medications,³⁻⁵ supportive therapy,^{6,7} client-centered therapy,⁸ meaning-oriented therapy,⁹ brief dynamic therapy,^{10,11} cognitive therapy,¹² cognitive behavioral therapy (CBT),¹³⁻¹⁷ interpersonal therapy (IPT),¹⁸ pastoral counseling,¹⁹ play therapy,²⁰ logotherapy,²¹ writing therapy,^{22,23} Internet-administered therapy,^{24,25} virtual reality,²⁶ and hypnosis.²⁷⁻²⁹ These treatments have been tested with children³⁰⁻³⁴ and adults^{10,22,35} and have included interventions for inpatients,³⁶ refugees,¹⁷ couples,³⁷ parents,³⁵ and those bereaved by war,³⁸ natural disasters,³⁹ accidents,²³ suicide,⁴⁰ and violence.⁴¹

Relatively few of these interventions have targeted complicated grief (CG) symptoms specifically rather than depression and distress more generally. Three review articles have described the literature on these CG-specific interventions.⁴²⁻⁴⁴ The most recent, a meta-analysis of ran-

Address for correspondence: Julie Wetherell, PhD, UCSD Department of Psychiatry, 9500 Gilman Drive, Dept. 9111N-1, La Jolla, CA 92093-9111, USA (e-mail: jwetherell@ucsd.edu)

Clinical research

domized, controlled trials, found a pooled standard mean difference (a measure of effect size) of -0.53 (95% CI: -1.00 to -0.07) favoring interventions targeting complicated grief relative to supportive counseling, IPT, or wait list.⁴³ The four interventions that were more efficacious than the comparison condition were all based, at least in part, on cognitive-behavioral principles.^{14,24,45} An interpretive intervention focused on increasing clients' insight about conflict and trauma related to their loss was not efficacious.^{46,47} The effects of the CG interventions appeared to grow larger at follow-up, although long-term data were only available from a single study.¹⁴

One form of complicated grief therapy (CGT) with strong empirical support has roots in both IPT and CBT.⁴⁵ CGT is based on attachment theory, which holds that humans are biologically programmed to seek, form, and maintain close relationships. Attachment figures are people with whom proximity is sought and separation resisted; they provide a "safe haven" of support and reassurance under stress and a "secure base" of support for autonomy and competence that facilitates exploration of the world. In acute grief following the loss of an attachment figure, the attachment system is disrupted, often leading to a sense of disbelief, painful emotions, intrusive thoughts of the deceased individual, and inhibition of the exploratory system.⁴⁸ With successful mourning, the individual moves from a state of acute grief to integrated grief in which the finality of the loss is acknowledged, the trauma of the loss is resolved, emotions become more positive or bittersweet, the mental representation is revised to encompass the death of the attachment figure, and the exploratory system is reactivated, with life goals revised to integrate the consequences of the loss. This occurs through a "dual-process model," with both loss- and restoration-focused activities.

In CG, the process of transition from acute grief to integrated grief is derailed.⁴⁹ Clients with CG typically experience prolonged, intense painful emotions; rumination, often around themes of self-blame; and maladaptive behaviors, including avoidance of triggers to the extent that functioning is disrupted.⁵⁰ Although the causes are not yet understood, the mechanism is believed to be incomplete processing of information about the death. Specifically, the mental representation of the attachment figure is disrupted, such that the loss is acknowledged in declarative memory but not in implicit memory. This leads to a lack of acceptance of the finality of the loss.

Moreover, the exploratory system does not re-engage, such that the grieving individual can become distanced from other people and the world generally.

Thus, the basic principle underlying CGT is that grief is a natural, adaptive process.⁵¹ This implies that treatment of CG involves removing the impediments to successful resolution of the grieving process. Through a variety of loss- and restoration-focused techniques, the therapist works to facilitate the progress of grief to help the client come to terms with the death.

A number of investigations have provided empirical support for this model of treatment. After initial pilot studies showed promising results,^{52,53} CGT was compared with standard IPT in a randomized trial with 83 adult outpatients with complicated grief.⁴⁵ Participants in both conditions received 16 individual sessions of psychotherapy. Treatment response was defined as a score of 1 or 2 ("very much improved" or "much improved") on the interviewer-rated Clinical Global Impression – Improvement scale and as time to a 20-point or better decrease in scores on the self-reported Inventory of Complicated Grief. Response rates were higher (51% vs 28%) and time to response faster in the CGT group than in the IPT group.

A secondary analysis examining the impact of naturalistic pharmacotherapy on participants in this trial found that response rates in the CGT group were higher among those taking antidepressant medications, and that this effect was mediated by reduced attrition among those taking medications.⁵⁴ Among patients receiving CGT, 42% of those not taking antidepressants, vs only 9% of those taking such medication, terminated the trial prematurely. By contrast, in the IPT condition, only 30% of those taking medications and 23% of those not taking medications dropped out. These data suggest that CGT may be a challenging treatment, particularly for individuals who are not also taking medication.

Investigators have subsequently tested CGT with Japanese women bereaved by violent death⁵⁵ and in substance abusers⁵⁶; results suggest that the benefits of treatment are not restricted to Western cultures or individuals without comorbid drug or alcohol abuse.

Description of the treatment

As noted above, the theory includes elements drawn from both IPT and CBT. In general, the CBT techniques target the loss-related processes and focus on symptoms

of painful intrusive memories and behavioral avoidance. The IPT elements focus on restoration by helping clients re-establish relationships and connection with valued life goals.

Although CGT can be flexibly applied in clinical practice, the manualized form tested in research studies consists of 16 sessions, each approximately 45 to 60 minutes long. Each session is structured, with an agenda that includes reviewing the previous week's activities, doing work in session, and assigning tasks for the coming week. The treatment is typically divided into three phases. In the introductory phase, which usually takes place over the first three sessions, the primary goals are to establish a strong therapeutic alliance, obtain a history of the client's interpersonal relationships, provide psychoeducation about the model of complicated grief, and describe the elements of treatment. A supportive person usually attends the third session. In the intermediate phase, which typically comprises sessions 4 to 9, the client performs a number of exercises inside and outside of the session designed to come to terms with the loss and address restoration of the capacity for joy and satisfaction in life. In the final sessions (10 to 16), the therapist and client review progress and collaboratively decide how to use the remaining sessions to complete the work and consolidate treatment gains. For some clients, this portion of the treatment may resemble IPT. A more detailed, session-by-session description follows.

Session 1

The goals of the first session are to welcome clients and orient them to CG and its treatment. Consistent with CGT's roots in interpersonal therapy, the primary focus of session 1 is to obtain an interpersonal history including early family relationships, other losses, the relationship with the deceased and the story of the death, and current relationships. The therapist and client discuss the client's current life situation, including stressors and coping resources. The therapist also provides a very brief introduction to the rationale and processes involved in CGT. Finally, the therapist introduces between-session assignments (sometimes known as homework): the grief monitoring diary, on which clients record daily triggers and less distressing moments; interval plans, which can include at-home practice of CG exercises as well as individualized activities designed to help clients move closer to their aspirations; and a handout that describes in

detail the model of CG and an overview of the treatment.

Session 2

In the second session, the therapist and client review the grief monitoring diary, examining triggers throughout the week and times when grief was relatively manageable to look for patterns. They also use the handout to discuss the model of CG and ways in which it relates to the client's situation. The therapist then provides an overview of the treatment. Finally, the client is encouraged to think about personal aspirations, activities that have the potential for reawakening the capacity for joy and meaning in life. The client is also given another copy of the CGT handout to provide to a supportive person who will attend the third session.

Session 3

Usually session 3 includes a supportive person such as a family member or close friend, either in person or, if necessary, by telephone. The rationales for including a supportive person are that individuals experiencing complicated grief often lose a sense of connection with others, which the treatment aims to help restore; an outside perspective on the client and the way that grief is affecting his or her life can be helpful for the therapist; and a friend or family member can facilitate the treatment by understanding what the client is doing and why, and providing support throughout the process, which is often difficult and painful. During the session, this individual is asked to describe the client since the death, his or her reactions to grief, and any avoided situations or activities. The therapist then provides an overview of the CG model and treatment to the support person. The client and support person discuss ways in which the latter can be helpful as the client progresses through the treatment. During the last 15 minutes or so, the client is seen alone to review the grief monitoring diary and provide an update on goal work.

Session 4

The heart of CGT begins in this session, with the introduction of imaginal revisiting. Imaginal revisiting is a core element of CGT that in some ways resembles prolonged exposure, an empirically supported therapy for

Clinical research

trauma and post-traumatic stress disorder (PTSD).^{57,58} In this technique, the client briefly (for approximately 5 minutes) visualizes and tells the story of when he or she became aware of the loved one's death into a tape recorder and then debriefs with the therapist. The goal of the exercise is to help the client come to terms with the loss by processing it at an emotional level and integrating that emotional processing with the rational knowledge that the loved one has died. In the debriefing portion of the exercise, the client describes what he or she observed while telling the story; the function of this discussion is to encourage the client to reflect on the story from the vantage point of the present. The client then participates in another visualization exercise in which the story is put away. Finally, clients identify a reward they can give themselves for doing the hard, painful work of revisiting, both in session and during the assignment of listening to the tape every day between sessions. Other elements that continue throughout the treatment include the grief monitoring diary and restoration-oriented work to help the client move toward a personal goal that is unrelated to grief, in order to begin to visualize life with the capacity for joy and satisfaction without the loved one who died.

Session 5

This session includes a review of the grief monitoring diary, imaginal revisiting, and restoration work. Situational revisiting is a new element introduced during this session, in which the client identifies activities or places previously avoided because they trigger grief or serve as reminders of the loved one. The client is encouraged to engage in a situational revisiting activity every day.

Sessions 6 to 9

In addition to reviewing the grief monitoring diary, imaginal and situational revisiting, and aspirations work, the client completes a series of forms identifying pleasant memories and positive aspects or characteristics of the person who died as well as unpleasant memories/less positive aspects. Clients usually bring photographs and other mementos to some of these sessions.

Session 10

In this session, the therapist uses one or more structured

questionnaires, such as the Inventory for Complicated Grief, to help the client evaluate progress and identify “stuck” points. Together, they decide on a direction for the remainder of the treatment. These can include work on other losses or IPT-oriented relationship work related to interpersonal disputes or role transitions.

Sessions 11 to 16

In these sessions, clients continue to complete grief monitoring diaries, situational revisiting exercises, and aspirations work. Although typically imaginal revisiting work is no longer necessary (as determined by distress ratings remaining low throughout the exercise), additional exercises may be conducted if needed. One final exercise that can be helpful in bringing a sense of closure and closeness with the deceased loved one is the imaginal conversation. In this exercise, the client imagines that the loved one has just died but is able to hear and speak. The client then engages in an imaginal conversation, playing both the role of the self and also of the loved one. During this conversation, the client can ask questions and, speaking as the dead person, can respond and/or offer reassurance. Although this exercise is optional (and best performed in cases in which the relationship was positive), it can be a moving and meaningful experience for clients.

If the client is experiencing CG from multiple losses, exercises such as imaginal and situational revisiting may be performed around another death. Usually the progress of therapy for treating other losses is faster after completion of the process for the initial, most distressing loss. Clients may also choose to engage in other work that is less directly related to CG and is usually consistent with the IPT targets of role transition or relationship conflict. Techniques can include standard IPT techniques such as close analysis of problematic interactions and role plays.

The final task of sessions 11 to 16 is termination with the therapist. For some clients, this is seen as a positive development, a “graduation” marking the progress from intense and debilitating grief to a sense of healing and wholeness. For other clients, discussion is required to process the feelings of loss of the therapeutic relationship.

Case example

The client, “Ann,” was a 52-year-old woman mourning the loss of her husband 4 years previously from a sudden

cardiac arrest. She had been abused in childhood, and the only truly satisfying relationship of her life had been with her husband, whom she met in her late 30s. She described him as a soul mate and best friend. They had chosen not to have children and in her words, “were everything to each other.” In addition to her emotional loss, her financial status deteriorated dramatically after his death, resulting in major life changes and a drop in her standard of living. She indicated that her husband had left their financial affairs in disarray, with records stored in boxes in the closet, but she did not feel capable of sorting through the boxes and dealing with the estate or taxes.

At the outset of therapy, Ann met criteria for major depression and PTSD as well as CG. She described crying every night, with great difficulty sleeping. She felt isolated from other people and did not socialize with former friends or colleagues at work. She stated that spending time with people she used to spend time with as part of a couple was too painful, and that although she knew it was irrational, she found herself feeling envious and resentful of other people’s relationships. She reported that she spent hours every day engaged in reveries about her life with her husband; her inattentiveness had drawn reprimands from her supervisor and she was now worried about losing her job, which she detested but needed for financial reasons. She had nightmares and flashbacks about waking to discover her husband’s body in their bed. She also reported episodes of rage, usually triggered by hearing about what she perceived as medical malpractice or instances of poor medical care. She held her husband’s physicians responsible for his death because they never diagnosed his heart problem. Although she had formerly been a talented amateur musician, playing guitar in a local band, she had not played since her husband’s death. She told the therapist that although she would never consider suicide, she could not imagine a future for herself without her husband.

Ann was able to complete the grief monitoring diary and rapidly recognized a pattern: although she disliked her job, work was a useful distraction from the pain she experienced as unrelenting in the evenings and on weekends. She found the description of CG in the handout reassuring, because it gave a name to her experiences. She saw many aspects of herself and her situation in the material. Upon hearing about the revisiting exercises, particularly the imaginal revisiting, she became very anx-

ious. She asked many questions about how telling the horrifying story of waking up in bed next to her deceased husband could possibly be helpful. In addition to explaining several times the ways in which this technique facilitates healthy grieving, the therapist also repeatedly reassured Ann that she would not be going through this experience alone.

Because she did not have any close friends who lived locally, she arranged to have her sister attend the third session by conference call. She was surprised at how supportive her sister was. The sister indicated that she was aware that Ann was suffering a great deal but had not known how to help and was “afraid to make things worse by saying the wrong thing.” She agreed to text Ann every day and talk with her twice a week, including the evening of her therapy appointments.

Ann initially had difficulty with imaginal revisiting. At the beginning of session 4, she asked a lot of questions about the rationale and procedures for the exercise; most of these were the same questions she had asked during session 2. The therapist normalized her concerns and praised her willingness to do something painful to help resolve her grief and come to terms with the loss. Because Ann was so hesitant to begin, the therapist also told her only to spend 2 minutes during the first exercise. Ann did so and was, as she expected, very distressed. During the debriefing process, Ann sobbed as she expressed her guilt over having slept through her husband’s passing and her agony at not knowing whether she could have saved him had she been awake. She also expressed anger toward her husband’s primary care doctor, who had performed routine annual physical examinations but had never diagnosed cardiac problems. She was able to perform the visualization exercise aimed at putting the story away and reported a decrease in distress to manageable levels. Although she agreed to listen to the tape between sessions and scheduled a telephone check-in with her therapist after completing the exercise the first time, when the time came, Ann told the therapist she was not yet ready.

After doing the imaginal revisiting exercise again in session 5, Ann reported that it was still very distressing, but she was willing to try listening to the tape at home. She and the therapist talked about ways Ann could reward herself for her hard work. She decided she would try to play her guitar, which had always been very pleasant. This time, she was able to complete the imaginal revisiting several times during the week and reported that

Clinical research

although it was painful, it was less hard than she had imagined it would be. Throughout the next 6 sessions, she continued to engage in exercises and spent most of the debriefing time focused on the issues of guilt, uncertainty, and anger, which Ann and the therapist agreed were the key factors contributing to her CG.

In contrast to the imaginal revisiting, Ann took a great deal of satisfaction from the situational revisiting, which began in session 5. She began by dining in a few of her husband's favorite restaurants that she had not visited since his death. She opted for lunches rather than dinners, invited acquaintances from work to accompany her, and found that rather than being distressed, she enjoyed both the food and the company. She then began the task of going through the files and dealing with the financial and tax issues. Although this was clearly not pleasant, Ann felt very proud of herself for taking responsibility and was able to make progress over the course of her treatment.

During session 2, Ann had identified playing in a band as one of her aspirations, and she began playing guitar again as a reward for doing the imaginal revisiting. She was soon back in her former routine of daily practice. After 2 months, she reported to her therapist that she had attended an informal jam session at a local bar and met a couple of musicians who were looking for a guitarist. They began practicing together and by the end of treatment had made plans to appear together as a band at the same bar for a performance that had the potential to turn into a monthly gig. Ann was delighted to be playing regularly and had begun to consider her new bandmates as friends.

Ann's ninth session was the week of what would have been her 10th wedding anniversary. The therapist began discussing this potentially painful time several weeks in advance. Two weeks before the anniversary, the therapist provided Ann with a handout on dealing with difficult times. With mixed feelings, Ann made plans to leave town for a long weekend to attend a music festival that she knew her late husband would not have enjoyed. Although she reported feeling very sad, she did enjoy the festival and felt relieved that she had not spent the time "moping" at home.

By the end of treatment, Ann had made a great deal of progress. She no longer met criteria for major depression, PTSD, or complicated grief. Although she continued to experience some moments and even days of sadness, these were not prolonged or incapacitating. She no longer experienced nightmares or flashbacks. She no

longer felt guilty and felt much less angry toward doctors and the medical profession. She no longer avoided pictures or places that reminded her of her husband. She now considered her bandmates and several of her coworkers as friends and had forged a closer relationship with her sister. She had also made progress on disentangling her financial affairs, although some of the issues were still unresolved. She was playing guitar in a band once again, which gave her great satisfaction.

Termination with the therapist was an easy process for Ann. She expressed gratitude for the help and accepted praise for her hard work, particularly in sticking with the imaginal revisiting exercise, which she was able to acknowledge helped reduce the power and pain of the memory of finding her husband's body. She agreed that she was doing much better and did not feel the need for further counseling at this time.

Current and future directions for research on complicated grief therapy

Research on CGT is ongoing. One currently unresolved issue is the role of pharmacotherapy in the recovery from CG. One investigation based on naturalistic data suggested that concurrent use of antidepressant medications may facilitate CGT by enabling clients to tolerate the painful work of imaginal and situational revisiting.⁵⁴ An open-label pilot study suggested that selective serotonin reuptake inhibitors may be sufficient to treat CG even in the absence of psychotherapy.⁵ Because CGT is a challenging treatment not yet widely available, a finding that medication alone is sufficient to alleviate suffering in many individuals would have important public health significance.

Currently, a large-scale trial is underway in four sites to investigate these questions. Clients with CG as indicated by a score of 30 or more on the Inventory of Complicated Grief⁵⁹ are randomly assigned to citalopram, pill placebo, CGT plus placebo, or CGT plus citalopram. The primary aim is to determine whether citalopram is more effective than placebo in reducing the symptoms of CG, as measured by the Clinician Global Impression - Improvement.⁶⁰

Another area ripe for exploration is the disseminability of CGT. Drawing as it does from both IPT and CBT, it can be challenging to learn for therapists who have a strong background in one model but not in the other. Like other therapies that deal with intense pain, it can

also be emotionally draining. To date, the process for obtaining the requisite skills to conduct CGT competently has involved a multi-day didactic workshop followed by intensive supervision of at least two cases, with an expert supervisor listening to audiotapes on an hour-for-hour basis. This level of training and supervision may

not be readily available for all potential therapists. It would be of interest to investigate whether a less stringent, time-intensive training process is sufficient to produce good outcomes; such a finding would greatly increase the public health significance of this promising new therapy. □

El tratamiento del duelo complicado como una nueva aproximación terapéutica

La terapia del duelo complicado (TDC) es un modelo de psicoterapia relativamente nuevo diseñado para manejar los síntomas del duelo complicado. La TDC está extraída de la teoría del apego y tiene raíces en la terapia interpersonal (TIP) y en la terapia cognitivo-conductual e incluye técnicas similares a la exposición prolongada (relatar repetidamente la historia de la muerte y realizar actividades de exposición en vivo). El tratamiento también incluye centrarse en los objetivos y en las relaciones personales. Se demostró la eficacia de la TDC en un ensayo en el que los participantes con duelo complicado fueron asignados aleatoriamente a la TDC o a la TIP, y los sujetos que recibieron TDC respondieron más rápidamente y con mejor respuesta global (51% v/s 28%). Este artículo resume brevemente las bases conceptuales de la TDC, discute la evidencia empírica para su eficacia, describe su técnica y presenta como ejemplo el caso de un cliente tratado con un protocolo manualizado de TDC de 16 sesiones. Se concluye con una descripción de las futuras líneas de investigación de la TDC.

Le traitement du deuil compliqué : une nouvelle approche

Le traitement du deuil compliqué (TDC) est un modèle relativement nouveau de psychothérapie destiné à en traiter les symptômes. Issu de la théorie de l'attachement et enraciné à la fois dans la thérapie interpersonnelle (TIP) et dans la thérapie cognitivocomportementale (TCC), le TDC comprend des techniques proches de l'exposition prolongée (récit répété de l'histoire de la mort et exposition in vivo). Le traitement s'intéresse aussi aux buts et aux relations personnelles. L'efficacité du TDC a été démontrée dans une étude dont les participants souffrant de DC ont été répartis de façon randomisée entre la TIP et le TDC ; ceux qui ont bénéficié du TDC ont répondu plus rapidement et ont été globalement plus nombreux à répondre (51 % vs 28 %). Cet article résume brièvement les concepts de base du TDC, analyse les preuves empiriques de son efficacité, décrit ses techniques et présente un cas clinique d'un patient traité selon un protocole de TDC en 16 séances. L'article conclut sur une description des futures directions de recherche pour le TDC.

REFERENCES

- Hensley PL. A review of bereavement-related depression and complicated grief. *Psychiatr Annals*. 2006;36:619-626.
- Hensley PL. Treatment of bereavement-related depression and traumatic grief. *J Affect Disord*. 2006;92:117-124.
- Hensley PL, Slonimski CK, Uhlenhuth EH, Clayton PJ. Escitalopram: an open-label study of bereavement-related depression and grief. *J Affect Disord*. 2009;113:142-149.
- Zygmunt M, Prigerson HG, Houck PR et al. A post hoc comparison of paroxetine and nortriptyline for symptoms of traumatic grief. *J Clin Psychiatry*. 1998;59:241-245.
- Simon NM, Thompson EH, Pollack MH, Shear MK. Complicated grief: a case series using escitalopram. *Am J Psychiatry*. 2007;164:1760-1761.
- Piper WE, Ogrodniczuk JS, Joyce AS, Weideman R. *Short-Term Group Therapies for Complicated Grief: Two Research-Based Models*. Washington, DC: American Psychological Association; 2011.
- Webb NB. Groups for children traumatically bereaved by the attacks of September 11, 2001. *Int J Group Psychother*. 2005;55:355-374.
- Goodman RF, Morgan AV, Juriga S, Brown EJ. Letting the story unfold: a case study of client-centered therapy for childhood traumatic grief. *Harv Rev Psychiatry*. 2004;12:199-212.
- Neimeyer RA, Wogrin C. Psychotherapy for complicated bereavement: a meaning-oriented approach. *Illness, Crisis, Loss*. 2008;16:1-20.
- Horowitz MJ, Marmar C, Weiss DS, DeWitt KN, Rosenbaum R. Brief psychotherapy of bereavement reactions. The relationship of process to outcome. *Arch Gen Psychiatry*. 1984;41:438-448.
- Marmar CR, Horowitz MJ, Weiss DS, Wilner NR, Kaltreider NB. A controlled trial of brief psychotherapy and mutual help group treatment of conjugal bereavement. *Am J Psychiatry*. 1988;145:203-209.
- Malkinson R. *Cognitive Grief Therapy: Constructing a Rational Meaning to Life Following Loss*. New York, NY: W. W. Norton; 2007.
- Allen B, Oseni A, Allen KE. The evidence-based treatment of chronic posttraumatic stress disorder and traumatic grief in an adolescent: a case study. *Psychol Trauma: Theory, Res, Prac, Policy*. In press.

Clinical research

14. Boelen PA, de Keijser J, van den Hout MA, van den Bout J. Treatment of complicated grief: a comparison between cognitive-behavioral therapy and supportive counseling. *J Consult Clin Psychology*. 2007;75:277-284.
15. Brown EJ, Pearlman MY, Goodman RF. Facing fears and sadness: cognitive-behavioral therapy for childhood traumatic grief. *Harv Rev Psychiatry*. 2004;12:187-198.
16. Cohen JA, Mannarino AP, Staron VR. A pilot study of modified cognitive-behavioral therapy for childhood traumatic grief (CBT-CTG). *J Am Acad Child Adolesc Psychiatry*. 2006;45:1465-1473.
17. Murray LK, Cohen JA, Ellis BH, Mannarino A. Cognitive behavioral therapy for symptoms of trauma and traumatic grief in refugee youth. *Child Adolesc Psychiatry Clin N Am*. 2008;17:585-604.
18. Crenshaw DA. An interpersonal neurobiological-informed treatment model for childhood complicated grief. *Omega*. 2006;54:319-335.
19. Blakley TL. Murder and faith: a reflected case study of pastoral interventions in traumatic grief. *J Pastoral Care Counsel*. 2007;61:59-69.
20. Carter R. Use of puppets to treat traumatic grief: a case study. *Elementary School Guidance Couns*. 1987;21:210-215.
21. Rogina JM, Quilitch HR. Logotherapy treatment of complicated grief syndrome. *Int Forum Logotherapy*. 2010;33:30-41.
22. O'Connor M, Nikolettis S, Kristjanson LJ, Loh R, Willcock B. Writing therapy for the bereaved: evaluation of an intervention. *J Palliat Med*. 2003;6:195-204.
23. Range LM, Kovac SH, Marion MS. Does writing about the bereavement lessen grief following sudden, unintentional death? *Death Stud*. 2000;24:115-134.
24. Wagner B, Knaevelsrud C, Maercker A. Internet-based cognitive-behavioral therapy for complicated grief: a randomized controlled trial. *Death Stud*. 2006;30:429-453.
25. Wagner B, Maercker A. A 1.5-year follow-up of an internet-based intervention for complicated grief. *J Trauma Stress*. 2007;20:625-629.
26. Botella C, Osma J, Palacios AGT, Guillén V, Baños R. Treatment of complicated grief using virtual reality: a case report. *Death Stud*. 2008;32:674-692.
27. Iglesias A, Iglesias A. Hypnotic treatment of PTSD in children who have complicated bereavement. *Am J Clin Hypn*. 2005;48:183-189.
28. van der Hart O, Brown P, Turco RN. Hypnotherapy for traumatic grief: Janetian and modern approaches integrated. *Am J Clin Hypn*. 1990;32:263-271.
29. Vijselaar J, van der Hart O. The first report of hypnotic treatment of traumatic grief: a brief communication. *Int J Clin Exp Hypn*. 1992;40:1-6.
30. Cohen JA, Mannarino AP. Treatment of childhood traumatic grief. *J Clin Child Adolesc Psychol*. 2004;33:819-831.
31. Cohen JA, Mannarino AP, Knudsen K. Treating childhood traumatic grief: a pilot study. *J Am Acad Child Adolesc Psychiatry*. 2004;43:1225-1233.
32. Kerig PK, Sink HE, Cuellar RE, Vanderzee KL, Elfstrom JL. Implementing trauma-focused CBT with fidelity and flexibility: a family case study. *J Clin Child Adolesc Psychol*. 2010;39:713-722.
33. McClatchey IS, Vonk ME, Palardy G. Efficacy of a camp-based intervention for childhood traumatic grief. *Res Soc Work Pract*. 2009;19:19-30.
34. Saltzman WR, Pynoos RS, Layne CM, Steinberg AM, Aisenberg E. Trauma- and grief-focused intervention for adolescents exposed to community violence: results of a school-based screening and group treatment protocol. *Group Dynamics*. 2001;5:291-303.
35. Oliver RC, Sturtevant JP, Schetz JP, Fallat ME. Beneficial effects of a hospital bereavement intervention program after traumatic childhood death. *J Trauma*. 2001;50:440-446.
36. Rosner R, Lumbeck G, Geissner E. Effectiveness of an inpatient group therapy for comorbid complicated grief disorder. *Psychother Res*. 2011;21:210-218.
37. Swan AH, Scott C. Complicated Grief: implications for the treatment of post-traumatic stress disorder in couples. *Sex Rel Ther* 2009;24:16-29.
38. Cox J, Davies DR, Burlingame GM, Campbell JE, Layne CM, Katzenbach RJ. Effectiveness of a trauma/grief-focused group intervention: a qualitative study with war-exposed Bosnian adolescents. *Int J Group Psychother*. 2007;57:319-345.
39. Salloum A, Overstreet S. Evaluation of individual and group grief and trauma interventions for children post disaster. *J Clin Child Adolesc Psychol*. 2008;37:495-507.
40. de Groot M, Neeleman J, van der Meer K, Burger H. The effectiveness of family-based cognitive-behavior grief therapy to prevent complicated grief in relatives of suicide victims: the mediating role of suicide ideation. *Suicide Life Threat Behav*. 2010;40:425-437.
41. Blakley, 2007; Miller L. Line-of-duty death: psychological treatment of traumatic bereavement in law enforcement. *Int J Emerg Ment Health*. 2007;9:13-23.
42. Jacobs S, Prigerson H. Psychotherapy of traumatic grief: a review of evidence for psychotherapeutic treatments. *Death Stud*. 2000;24:479-495.
43. Wittouck C, Van Autreve S, De Jaegere E, Portzky G, van Heeringen K. The prevention and treatment of complicated grief: a meta-analysis. *Clin Psychol Rev*. 2011;31:69-78.
44. Zhang B, El-Jawahri A, Prigerson HG. Update on bereavement research: Evidence-based guidelines for the diagnosis and treatment of complicated bereavement. *J Palliat Med*. 2006;9:1188-1203.
45. Shear MK, Frank E, Houck PR, Reynolds CF. Treatment of complicated grief. A randomized controlled trial. *JAMA*. 2005;293:2601-2608.
46. Piper WE, McCallum M, Joyce AS, Rosie JS, Ogrodniczuk JS. Patient personality and time-limited group psychotherapy for complicated grief. *Int J Group Psychotherapy*. 2001;51:525-552.
47. Piper WE, Ogrodniczuk JS, Joyce AS, Weideman R, Rosie JS. Group composition and group therapy for complicated grief. *J Consult Clin Psychology*. 2007;75:116-125.
48. Shear K, Shair H. Attachment, loss, and complicated grief. *Dev Psychobiol*. 2005;47:253-267.
49. Shear MK, Zuckoff A, Frank E. The syndrome of traumatic grief. *CNS Spectr*. 2001;6:339-346.
50. Shear MK. Exploring the role of experiential avoidance from the perspective of attachment theory and the dual process model. *Omega*. 2010;61:357-369.
51. Shear MK. Complicated grief treatment: the theory, practice, and outcomes. *Bereave Care*. 2010;29:10-14.
52. Harkness KL, Shear MK, Frank E, Silberman RA. Traumatic grief treatment: case histories of 4 patients. *J Clin Psychiatry*. 2002;63:1113-1120.
53. Shear MK, Frank E, Foa E, et al. Traumatic grief treatment: a pilot study. *Am J Psychiatry*. 2001;158:1506-1508.
54. Simon NM, Shear MK, Fagiolini A et al. Impact of concurrent naturalistic pharmacotherapy on psychotherapy of complicated grief. *Psychiatry Res*. 2008;159:31-36.
55. Asukai N, Tsuruta N, Saito A. Pilot study on traumatic grief treatment program for Japanese women bereaved by violent death. *J Trauma Stress*. 2011;24:470-473.
56. Zuckoff A, Shear K, Frank E, Daley DC, Seligman K, Silowash R. Treating complicated grief and substance use disorders: a pilot study. *J Subst Abuse Treat*. 2006;30:205-211.
57. Foa EB, Rothbaum BO, Riggs DS, Murdock TB. Treatment of posttraumatic stress disorder in rape victims: a comparison between cognitive-behavioral procedures and counseling. *J Consult Clin Psychol*. 1991;59:715-723.
58. Powers MB, Halpern JM, Ferenschak MP, Gillihan SJ, Foa EB. A meta-analytic review of prolonged exposure for posttraumatic stress disorder. *Clin Psychol Rev*. 2010;30:635-641.
59. Prigerson HG, Maciejewski PK, Reynolds CF 3rd, et al. Inventory of Complicated Grief: a scale to measure maladaptive symptoms of loss. *Psychiatry Res*. 1995;59:65-79.
60. Guy W. *ECDEU Assessment Manual for Psychopharmacology — Revised* (DHEW Publ No ADM 76-338). Rockville, MD: U.S. Department of Health, Education, and Welfare, Public Health Service, Alcohol, Drug Abuse, and Mental Health Administration, NIMH Psychopharmacology Research Branch, Division of Extramural Research Programs; 1976:218-222.